




One of the most worrisome impacts of climate change is a possible acceleration of the global water cycle leading to more frequent droughts and floods.


IMAGE 4




The ocean retains a better record of changes in precipitation than the land, and translates these changes into variations in the salt concentration of its surface waters.











The Research Vessel Knorr will serve as a platform for temperature, salinity, and meteorological measurements in addition to transporting a large fleet of robotic vehicles to the SPURS site in the tropical Atlantic.




Researchers will deploy an array of three oceanographic moorings at the SPURS site for a year: the WHOI surface flux mooring will measure meteorological parameters and fluxes between the ocean and atmosphere, while two NOAA profiling crawler ("prawler") moorings will obtain meteorological observations and serial profiles along the mooring wire.




Satellite-tracked surface drifters will track water movement. SPURS will deploy profiling floats, modified from those used in the global Argo program to measure near-surface salinity, wind speed, and rain rate, and Lagrangian (free-floating) drifters equipped with multiple sensors to accurately track three-dimensional water movement.



SPURS will use several types of autonomous underwater vehicles. Two propeller-driven IVER2/Ecomapper vehicles will conduct short-term and small-scale surveys of salinity features. Two buoyancy-driven Slocum Gliders equipped with microstructure sensors will observe the intensity of ocean mixing. Three high-endurance, buoyancy-driven Seagliders will measure temperature, salinity, and microstructure in a 93-mi (150-km) box surrounding the SPURS moorings for a year.




Three sun-powered wave gliders equipped to measure temperature, salinity, and meteorological conditions will occupy a 62-mi (100-km) region near the SPURS moored array for a year.




SPURS will use a nested observational strategy: expendable devices and ships-of-opportunity will collect temperature and salinity measurements over a wide geographic area in the central tropical Atlantic. Moorings and small fleets of robotic vehicles will provide a sustained year-long presence in a smaller observational area. During research cruises, ship-deployed platforms will make high-resolution measurements of even smaller oceanographic features.

Eric Lindstrom






NASA's Aquarius instrument has been collecting ocean surface salinity measurements from space since August 2011 and using data from Argo, a global array of thousands of drifting profiling floats measuring the temperature and salinity of the upper layers of the ocean, for comparison and validation.



The data collected by the SPURS devices will be relayed via satellite to shore and integrated with numerical ocean models and satellite observations.



Page Last Updated: September 5, 2012 Page Editor: Robert Garner NASA Official: Brian Dunbar NASA Information on the American Recovery and Reinvestment Act of 2009

Budgets, Strategic Plans and Accountability Reports Equal Employment Opportunity Data Posted Pursuant to the No Fear Act Information-Dissemination Policies and Inventories Freedom of Information Act Privacy Policy & Important Notices NASA Advisory Council Aerospace Safety Advisory Panel Inspector General Hotline Office of the Inspector General NASA Communications Policy Contact NASA Site Map BusinessUSA USA.gov Open Government at NASA Help and Preferences